

ORIGINAL RESEARCH PAPER

**SYNTHESIS OF GIBBERELLINS BY SURFACTANT
PRODUCERS *NOCARDIA VACCINII* IMV B-7405,
ACINETOBACTER CALCOACETICUS IMV B-7241 AND
RHODOCOCCUS ERYTHROPOLIS IMV AC-5017**

**Natalia Leonova¹, Tetyana Pirog^{1,2}, Daria Piatetska^{2*}, Tetyana
Shevchuk¹, Maxym Kharkhota¹, Galyna Iutynska¹**

¹*D.K. Zabolotny Institute of Microbiology and Virology of National Academy of
Science of Ukraine, 03143, Acad. Zabolotny str., 154, Kyiv, Ukraine*

²*National University of Food Technologies, Faculty of Biotechnology and
Ecological Control, 01601 Volodymyrska str., 68, Kyiv, Ukraine*

*Corresponding author: dashka2310@gmail.com

Received: February, 07, 2020

Accepted: November, 04, 2020

Abstract: In this study, we established that the producers of surfactants *Nocardia vaccinii* IMV B-7405, *Acinetobacter calcoaceticus* IMV B-7241 and *Rhodococcus erythropolis* IMV Ac-5017 were able to synthesize phytohormones of gibberellic nature during cultivation on the traditional substrates (ethanol, hexadecane) and on the industrial waste sunflower oil and biodiesel production waste). The results of specific biotesting showed that the treatment of cucumber seedlings with phytohormonal extracts of all studied strains at 1:500 and 1:600 dilutions stimulated the extension of the cucumber hypocotyls as compared to the water control in values close to the treatment with gibberellic acid. The results of high-performance liquid chromatography showed *N. vaccinii* IMV B-7405, *A. calcoaceticus* IMV B-7241 and *R. erythropolis* IMV Ac-5017 produce highly active forms of gibberellins GA₃ and GA₄. The level of its synthesis was nearly the same (6.0-10.0 μg·L⁻¹) under cultivation of strains on every substrate. The exception was strain *N. vaccinii* IMV B-7405 which synthesized almost 47.0 μg·L⁻¹ GA₃ and GA₄ while is growing on the waste oil from meat frying. The obtained results are the groundwork for the development of an economically profitable technology for the recycling of toxic wastes using *N. vaccinii* IMV B-7405, *A. calcoaceticus* IMV B-7241 and *R. erythropolis* IMV Ac-5017. Such technology will allow us to develop complex microbial preparations with various biological properties in a single process.

Keywords: *bacteria, GA₃, GA₄, phytohormones, specific biotesting, surface-active substances, toxic waste*

INTRODUCTION

Most researchers adhere to commonly accepted concept of “one producer – one product” while developing microbial biotechnologies and focusing their attention only on synthesis of the main target product. At the same time, the data about synthesis by microorganisms of several practically valuable metabolites, such as surface-active substances (surfactants) with polysaccharides, bacteriocins, polyhydroxyalkanoates or enzymes began to appear in the literature in the past few years [1 – 6]. The authors of these works note that the ability of strains to synthesize complexes of metabolites with various biological properties significantly expands the scope of their practical use.

We established the ability of *Nocardia vaccinii* IMV B-7405, *Acinetobacter calcoaceticus* IMV B-7241 and *Rhodococcus erythropolis* IMV Ac-5017 to synthesize exometabolites with phytohormonal activity (auxins and cytokinins) in the previous work [7]. We must note that synthesis of auxins and cytokinins depended on the nature of the source of carbon in the culture medium.

At the time of publication of the work [7], there were no data in the available literature on the ability of the surfactant producers to synthesize phytohormones. The production of indole-3-acetic acid by bacteria (mainly representatives of the genus *Rhodococcus*) isolated from soils contaminated with hydrocarbons and heavy metals was reported in 2016. But the authors were determining the surfactant synthesis ability by the emulsification index and reduction of surface tension, which was insignificant – no more than 60.0-65.0 mN·m⁻¹ (comparing to 30.0–35.0 mN·m⁻¹ by our producers of surfactants).

However, in recent years more and more publications have appeared about the microorganisms (mainly rhizobacteria) production of complexes of biologically active substances, in particular phytohormones and metabolites with antimicrobial, nematicidal, etc. activity [9 – 15]. Actinobacteria (in particular *Streptomyces*) [10 – 14], as well as the representatives of the genera *Bacillus* and *Paenibacillus* [9, 15], are the most active producers of complexes of such compounds with diverse biological activity. The analysis of the literature data [16 – 19] showed that, usually, microorganisms form phytohormones of the three classes - auxins, cytokinins and gibberellins, and this is true for those who associates with the plants (rhizospheric, endophytic, nitrogen-fixing, phytopathogenic bacteria), and for those who do not participate in such interaction.

The aim of the present work is to study the ability to synthesize compounds of gibberellic nature by surfactant producers *N. vaccinii* IMV B-7405, *A. calcoaceticus* IMV B-7241 and *R. erythropolis* IMV Ac-5017.

MATERIALS AND METHODS

Object of research

The objects of research are *Nocardia vaccinii* K-8, *Acinetobacter calcoaceticus* K-4 and *Rhodococcus erythropolis* EK-1 strains, registered in Microorganisms Depository of Institute of Microbiology and Virology, the National Academy of Sciences of Ukraine under the numbers IMV B-7405 (K-8), IMV B-7241 (K-4) and IMV Ac-5017(EK-1).

Medium composition and conditions of cultivation

N. vaccinii IMV B-7405 was grown in the liquid mineral medium ($\text{g}\cdot\text{L}^{-1}$ distilled water): $\text{NaNO}_3 - 0.5$, $\text{MgSO}_4\cdot 7\text{H}_2\text{O} - 0.1$, $\text{CaCl}_2\cdot 2\text{H}_2\text{O} - 0.1$, $\text{KH}_2\text{PO}_4 - 0.1$, $\text{FeSO}_4\cdot 7\text{H}_2\text{O} - 0.01$, yeast autolysate – 0.5 % v/v, pH 6.8–7.0.

Strain *A. calcoaceticus* IMV B-7241 was cultivated in the liquid medium ($\text{g}\cdot\text{L}^{-1}$ distilled water): $(\text{NH}_2)_2\text{CO}_3 - 0.35$, $\text{MgSO}_4\cdot 7\text{H}_2\text{O} - 0.1$, $\text{NaCl} - 1.0$, $\text{Na}_2\text{HPO}_4 - 0.6$, $\text{KH}_2\text{PO}_4 - 0.14$, pH 6.8–7.0. Yeast autolysate – 0.5 % v/v and solution of trace elements – 0.1 % v/v were also added to the medium. Trace elements solution contained ($\text{g}/100 \text{ mL}$): $\text{ZnSO}_4\cdot 7\text{H}_2\text{O} - 1.1$, $\text{MnSO}_4\cdot \text{H}_2\text{O} - 0.6$, $\text{FeSO}_4\cdot 7\text{H}_2\text{O} - 0.1$, $\text{CuSO}_4\cdot 5\text{H}_2\text{O} - 0.004$, $\text{CoSO}_4\cdot 7\text{H}_2\text{O} - 0.03$, $\text{H}_3\text{BO}_3 - 0.006$, $\text{KI} - 0.0001$, $\text{EDTA} - 0.5$.

Strain *R. erythropolis* IMV Ac-5017 was grown in the liquid mineral medium ($\text{g}\cdot\text{L}^{-1}$ distilled water): $\text{NaNO}_3 - 1.3$, $\text{NaCl} - 1.0$, $\text{Na}_2\text{HPO}_4\cdot 12\text{H}_2\text{O} - 0.6$, $\text{KH}_2\text{PO}_4 - 0.14$, $\text{MgSO}_4\cdot 7\text{H}_2\text{O} - 0.1$, $\text{FeSO}_4\cdot 7\text{H}_2\text{O} - 0.001$, pH 6.8–7.0.

The refined and fried sunflower oil (McDonald's Restaurant Network, Kiev, Ukraine), technical glycerol (Komsomolsk biofuel plant, Poltava region, Ukraine), *n*-hexadecane and ethanol were used as the carbon and energy sources in concentration of 2.0 % v/v.

The culture in the exponential phase was used as inoculum and added in concentration of 5 - 10 % of nutritive medium volume. The concentration of the corresponding carbon source in the medium for the inoculum obtainment was 0.5 % v/v.

The cultivation was carried out in 750 mL flasks, containing 100 mL of medium, on the shaker (320 rpm) at 28 – 30 °C during 7 days.

Determination of indexes of growth and biosurfactant synthesis

The biomass was determined by the gravimetric method. The double treatment of the culture broth with hexane or petrol ether was carried out to remove the residual *n*-hexadecane or oil respectively before the cell sedimentation (5000 g, 30 min). The necessity of this operation was caused by the precipitation of *n*-hexadecane or oil with cells that overestimate the biomass level.

The amount of exocellular surfactant was defined by weighing after it had been extracted from the culture broth supernatant with the mixture of chloroform and methanol (2 : 1). To obtain the supernatant, the culture broth was centrifuged at 5000 g for 20 minutes. Five milliliters of 1 N HCl were added to a 100 mL cylindrical separating funnel with 25 mL supernatant; the funnel was closed with the lap stopper and shaken for 3 minutes; then, a 15 mL mixture of chloroform and methanol (2 : 1) was added and stirred (extraction of lipids) for 5 minutes. The mixture obtained after the extraction was kept in the funnel until the separation of phases, then the lower fraction was poured (organic extract 1) and the water phase was re-extracted. At the repeated extraction, 5 mL of 1 N HCl solution and 15 mL of the mix of chloroform and methanol (2 : 1) were added to the water phase for extraction of lipids, which took 5 minutes. After the separation of the phases, the lower fraction was poured off and organic extract 2 was obtained. At the third step, 25 mL of chloroform and methanol (2 : 1) were added to the water phase and extraction was carried out as above. As a result, organic extract 3 was obtained. Extracts one to three were merged and evaporated in the IR-1M2 rotor evaporator (OJSC Khimlaborpribor, Russia) at 50 °C and at an absolute pressure of 0.4 atm until reaching constant mass.

Obtaining of extracts with phytohormonal activity (phytohormonal extracts)

After cultivation of the strains *N. vaccinii* IMV B-7405, *A. calcoaceticus* IMV B-7241 and *R. erythropolis* IMV Ac-5017, the biomass was separated by centrifugation (5000 g) for 25 min. Residuals of sunflower oil and hexadecane were extracted from the culture broth using petroleum ether or hexane (ratio 1 : 1).

Gibberellic extracts were obtained from the supernatant of the culture broth by extraction with ethyl acetate at pH 2.5. The extracts were evaporated to dryness under vacuum and redissolved in 80 % ethanol. The obtained extracts were stored at -24 °C. In the article, these preparations are called phytohormonal extracts.

Determination of gibberellic activity

To determine gibberellic activity we used hypocotyls of seedlings of cucumber grade Nezhinskiye by technique Brian's and Lemming's bioassay in the modification Ahnistrykova [21]. After germination of cucumber seeds during 3-4 days (temperature 27 °C) hypocotyls with length 2.0 ± 0.2 cm were selected. Hypocotyls were placed in Petri dishes with aliquots of aqueous solutions of the phytohormonal extracts at dilutions 1 : 400, 1 : 500 and 1 : 600. After incubation for 1 - 2 days at 27 °C elongation of hypocotyls was measured and it was compared to controls (treatment with distilled water and a solution gibberellic acid (GA₃) at a concentration of 10^{-5} M).

Qualitative and quantitative determination of gibberellins

Purification and concentration of the phytohormonal extracts were carried out on silicagel plates of the mark Silufol UV254 (Chemapol, Czech Republic) in a mixture of solvents used sequentially: chloroform, 12.5 % aqueous ammonia, ethyl acetate : acetic acid (20 : 1).

The qualitative and quantitative composition of gibberellins was analyzed by high performance liquid chromatography (HPLC) using the Agilent 1200 liquid chromatograph (Agilent Technologies, USA) and the Agilent G1956B mass spectrometry (MS) detector. The separation was carried out on analytical column Zorbax SB-C18 (2.1 mm × 150 mm, 3 μm; Agilent Technologies, USA), the flow rate of the mobile phase was $0.35 \text{ mL} \cdot \text{min}^{-1}$. The volumes of injections were 3 μL. Columns were kept at 30 °C. Elution was carried out in gradient mode in acetonitrile (A) - 0.1 % / water + formic acid (B): 20 % A for 5 minutes, then the content of A was gradually changed from 20 to 80 % for 10 minutes, followed by an increase of A to 100 % for 0.5 minutes. This ratio was kept for the next 8.5 minutes. Standard solutions of gibberellins GA₃, GA₄ and GA₇ (Sigma-Aldrich, Germany) were used for identification.

Detection of gibberellins was carried out using a diode-matrix detector with a signal recording at a wavelength of 198 and 210 nm. Mass spectrometric analysis was performed with the recording of positive and negative ions in the ratio m/z in the range 190 - 400 nm. The fluorescence detector was used in the extinction mode at 210 nm, and in the emission mode at 410 nm.

Analysis of the molecular weight of gibberellins

The molecular mass was determined using a single quadrupole mass spectrometric detector. Ionization was carried out in the mode of electrostatic spraying (ESI) with the formation of negative ions. Ion detection was performed in SCAN and SIM mode (selected ion monitoring) in the range of 200 ÷ 500 m/z. GA₃, GA₄ and GA₇ were registered by comparing the retention time, the values of the molecular weights of the ions, the spectral characteristics of the obtained peaks. The quantitative content of GA₃, GA₄ and GA₇ was determined by the external calibration method using the SIM mode by the 345, 331 and 329 m/z ions.

HPLC/MS analysis of gibberellic extracts of *N. vaccinii* IMV B-7405, *A. calcoaceticus* IMV B-7241 and *R. erythropolis* IMV Ac-5017 was made at the Center for collective use at the D.K. Zabolotny Institute of Microbiology and Virology of the NAS of Ukraine.

All the experiments were repeated three times, and the number of parallel measurements in each experiment made up 3 – 5. The statistical processing of the experimental data was carried out in accordance with the algorithm described in [22]. Differences of mean indicators were deemed as reliable at the significance level $p < 0.05$. All the used reagents were analytical grade.

RESULTS AND DISCUSSION

Synthesis of surfactants by *N. vaccinii* IMV B-7405, *A. calcoaceticus* IMV B-7241 and *R. erythropolis* IMV Ac-5017 on different substrates

The data of Table 1 confirm that the concentration of the surfactants synthesized by studied strains depended on nature of carbon source in cultivation medium. The highest synthesis rates were observed in case of using waste sunflower oil (2.3-3.5 g·L⁻¹) or technical glycerol (4.2 g·L⁻¹) as a substrate. The concentration of surfactants, synthesized on the traditional substrates (ethanol, hexadecane, refined oil) was lower (1.5-2.0 g·L⁻¹).

Table 1. The effect of nature of carbon source in culture medium of *N. vaccinii* IMV B-7405, *A. calcoaceticus* IMV B-7241 and *R. erythropolis* IMV Ac-5017 on synthesis of surfactants

| Strain | Carbon source in culture medium | Concentration of surfactant, [g·L ⁻¹] |
|------------------------------------|---------------------------------|---|
| <i>N. vaccinii</i> IMV B-7405 | Refined oil | 1.8±0.09 |
| | Waste oil after frying potatoes | 2.3±0.11 |
| | Waste oil after frying meat | 2.6±0.13 |
| <i>A. calcoaceticus</i> IMV B-7241 | Ethanol | 1.5±0.08 |
| | Technical glycerol | 4.2±0.21 |
| | Refined oil | 2.0±0.10 |
| | Waste oil after frying meat | 3.5±0.17 |
| <i>R. erythropolis</i> IMV Ac-5017 | Ethanol | 1.8±0.14 |
| | <i>n</i> -Hexadecane | 1.5±0.08 |
| | Refined oil | 1.8±0.09 |
| | Waste oil after frying meat | 2.3±0.11 |

Gibberellin-specific phytotesting

One of the approaches for studying the ability of gibberellin synthesis by bacteria is the phytotesting which gives us the opportunity to evaluate the biological effect of the phytohormonal extracts. Therefore, at the next stage, bioassay was performed using cucumber seedlings which are a typical test culture in gibberellin-specific phytotests.

Table 2. The effect of phytohormonal extracts of IMV B-7405, IMV B-7241 and IMV Ac-5017 on prolongation of hypocotyls of cucumbers grade Nezhinskiye

| Strain | Carbon source in culture medium | Length of hypocotyls (cm) after treatment by | | | |
|------------------------------------|---------------------------------|--|------------|-----------------|----------------------------|
| | | phytohormonal extracts in dilution | | water (control) | gibberellic acid (control) |
| | | 1 : 500 | 1 : 600 | | |
| <i>N. vaccinii</i> IMV B-7405 | Refined oil | 8.1 ± 0.1 | 8.4 ± 0.1 | 7.5 ± 0.1 | 8.3 ± 0.2 |
| | Waste oil after frying potatoes | 8.2 ± 0.1 | 8.1 ± 0.1 | 7.5 ± 0.1 | 8.4 ± 0.2 |
| | Waste oil after frying meat | 8.2 ± 0.1 | 8.5 ± 0.2 | 7.6 ± 0.1 | 8.5 ± 0.2 |
| <i>A. calcoaceticus</i> IMV B-7241 | Ethanol | 4.7 ± 0.1 | 4.8 ± 0.1 | 4.2 ± 0.1 | 4.6 ± 0.2 |
| | Technical glycerol | 5.0 ± 0.1 | 5.3 ± 0.1 | 4.7 ± 0.1 | 5.8 ± 0.3 |
| | Refined oil | 11.1 ± 0.2 | 11.3 ± 0.2 | 10.2 ± 0.1 | 11.4 ± 0.3 |
| | Waste oil after frying meat | 9.5 ± 0.1 | 9.1 ± 0.1 | 8.7 ± 0.1 | 9.8 ± 0.2 |
| <i>R. erythropolis</i> IMV Ac-5017 | Ethanol | 5.6 ± 0.2 | 6.0 ± 0.3 | 4.0 ± 0.1 | 5.9 ± 0.2 |
| | <i>n</i> -Hexadecane | 5.7 ± 0.1 | 5.8 ± 0.1 | 5.4 ± 0.1 | 6.0 ± 0.1 |
| | Refined oil | 6.5 ± 0.1 | 6.8 ± 0.2 | 5.7 ± 0.1 | 6.6 ± 0.2 |
| | Waste oil after frying meat | 7.2 ± 0.2 | 6.9 ± 0.1 | 5.8 ± 0.2 | 6.6 ± 0.2 |

Our studies have shown that in case of treatment of seedlings with phytohormonal extracts of strains *N. vaccinii* IMV B-7405, *A. calcoaceticus* IMV B-7241 and *R. erythropolis* IMV Ac-5017 the stimulation of prolongation of hypocotyls was observed comparing with using water or gibberellic acid solution for treatment (Table 2).

Data presented in Table 2 show that, regardless of the dilution of the extracts (500 or 600 times), the results of treatment were almost the same. However, with further 800 and 1000 times dilution the stimulation of hypocotyl prolongation wasn't observed. It should be noted that the effectiveness of the phytohormonal extracts depended on the nature of carbon source in cultivation medium of the surfactant producing strains.

Results obtained in the case of treatment of cucumber seedlings with the phytohormonal extracts in dilution 1 : 500, 1 : 600 are close to values obtained in the case of treatment with gibberellic acid. This allows us to assume that the extracts contain compounds of gibberellic nature.

The synthesis of gibberellins by *N. vaccinii* IMV B-7405, *A. calcoaceticus* IMV B-7241 and *R. erythropolis* IMV Ac-5017 on different substrates

Qualitative and quantitative composition of gibberellins in phytohormonal extracts of *N. vaccinii* IMV B-7405, *A. calcoaceticus* IMV B-7241 and *R. erythropolis* IMV Ac-5017 are presented in Table 3.

These data show that all studied strains are able to synthesize gibberellins A₃ and A₄, but the level of synthesis was different and depended on the nature of carbon source in the cultivation medium.




Figure 1. HPLC/MS analysis of metabolites of gibberellic nature extracted from the culture liquid supernatant of *N. vaccinii* IMV B-7405 under cultivation on the waste oil after frying potatoes. Shown are (A) GA₃ analysis and (B) GA₄ analysis

The highest concentration of GA₃ and gibberellin-synthesizing ability was observed for strain *N. vaccinii* IMV B-7405 under cultivation on the waste oil after frying potatoes or meat (Table 3, Figure 1). For other strains the rates of synthesis were an order of

magnitude lower. Under cultivation of *A. calcoaceticus* IMV B-7241 on the technical glycerol and *R. erythropolis* IMV Ac-5017 on the hexadecane the synthesis of gibberellic acid was almost not observed.

The data presented in Table 3 show that regardless of nature of carbon source in cultivation medium every strain had almost the same level of GA₄ synthesis: the concentration of GA₄ was in the range 5.63 -7.36 µg·L⁻¹, and gibberellin-synthesizing ability was within 3.77 - 8.38 µg GA₄/g of biomass.

Experiments have shown that all strains synthesized trace amounts of GA₇ in the studied conditions of cultivation.

Thus, the HPLC data are consistent with the phytotests data and indicate the ability of surfactants producing strains *N. vaccinii* IMV B-7405, *A. calcoaceticus* IMV B-7241 and *R. erythropolis* IMV Ac-5017 to synthesize GA₃, GA₄ and GA₇.

To date, the ability to synthesize phytohormones of gibberellic nature is established for a wide range of living organisms. Plants, bacteria, fungi and yeasts synthesize over 130 different forms of gibberellins [23]. However, only some of them demonstrate the high biological activity - its GA₁, GA₄, GA₄ and GA₇, and the other gibberellins are physiologically inactive and are the precursors of the biosynthesis of other forms.

Gibberellins are synthesized by many micromycetes [24 – 27], and not only by representatives of the genus *Fusarium* (*Gibberella*), which are the industrial producers of the gibberellic acid [28]. The synthesis of gibberellins by the endophytic fungi of the genus *Penicillium* is one of the mechanisms that allow the plants survive under salt stress [26]. As for the connection between pathogenicity and phytohormonal activity, the research of growth stimulating and pathogenic strains of *Fusarium culmorum* showed that the latter synthesized 4 times less gibberellin [27].

Many bacteria also synthesize gibberellins, moreover as phytopathogenic and associated with plants [18, 19, 29] and freely existing [30]. In the 80's of the 20th century was carried out the study of 10 wild and mutant (*nod*, *fix*) strains of *Rhizobium phaseoli* that showed that incapable of nodulation and nitrogen fixation mutants also synthesized gibberellins, and therefore, nitrogen-fixing ability is not related to phytohormonal activity [31].

Despite the fact that GA₄ and GA₇ are not inferior to GA₃ on biological activity, the industrial production for them is not developed. This is due to the following reasons: they are synthesized in extremely low concentrations, which causes high price of the preparations; as a rule, they are synthesized as a mixture of GA₄ and GA₇, in which the ratio GA₄/GA₇ varies greatly; excretion of individual preparations GA₄ and GA₇ from the mixture is complicated because of their very close polarity [32].

The ability to synthesize GA₄ and GA₇ has been identified as in microorganisms associated with plants - rhizospheric [33], endophytic [34], and in freely existing [35].

In 2017, a strain *Bacillus aryabhatai* SRB02 was isolated from soybean rhizosphere, which contributed to the better heat resistance of soybean plants. It produced the entire spectrum of biologically active gibberellins, among which prevailed GA₄ - 31 ng·L⁻¹, and the level of synthesis of others (GA₁ - 1 ng·L⁻¹, GA₃ - 6 ng·L⁻¹, GA₇ - 12 ng·L⁻¹) was lower [32].

Bacterial endophyte *Sphingomonas* sp. LK11 due to the ability to synthesize indol-3-acetic acid and gibberellins stimulates the growth of tomatoes (*Tephrosia apollinea*). In the culture broth of *Sphingomonas* sp. LK11, the presence of physiologically active GA₄ in the amount of 2970 ng·L⁻¹, as well as inactive GA₉ (980 ng·L⁻¹) and GA₂₀

(2410 ng·L⁻¹), was detected [34].

Strain *Leifsonia soli* sp. SE134 in the terms of experiment promoted the growth of cucumbers, tomatoes and young radishes, by increasing shoot length, plant fresh weight and chlorophyll in the leaves as compared with control. The wide spectrum of gibberellins, including GA₄ (15.8 ng·L⁻¹) and GA₇ (5.4 ng·L⁻¹), was identified in the culture broth of *L. soli* sp. SE134 [35].

Despite the fact that microorganisms of various physiological groups are able to synthesize gibberellins [23 – 35], in the available literature we could not find the information on the synthesis of gibberellins by the producers of surfactants.

Our research showed that producers of surfactants *N. vaccinii* IMV B-7405, *A. calcoaceticus* IMV B-7241 and *R. erythropolis* IMV Ac-5017 synthesize gibberellins GA₃, GA₄ and GA₇ under cultivation on the different substrates, including the industrial waste. It is worth mentioning that the level of gibberellin synthesis practically independent of the nature of the carbon source in the cultivation medium of the studied strains. An exception has appeared to be strain *N. vaccinii* IMV B-7405, which synthesized on the orders of magnitude higher amounts of GA₃ on the waste oil than on the refined oil. In our opinion it's because in the waste oil may be compounds, which are activators of enzymes of the gibberellin synthesis. Our further research will be devoted to clarifying this issue.

In Table 4 is presented generalized data on synthesis of the phytohormones of the three main classes - auxins, cytokinins and gibberellins, by the studied producers of surfactants.

Table 4. Influence of cultivation conditions of *N. vaccinii* IMV B-7405, *A. calcoaceticus* IMV B-7241 and *R. erythropolis* IMV Ac-5017 on synthesis of phytohormones

| Carbon source in culture medium | Strain | Concentration [µg·L ⁻¹] | | | |
|---------------------------------|-------------|-------------------------------------|----------------|--------------|---------------|
| | | auxins [2] | cytokinins [2] | gibberellins | Total |
| Ethanol | IMV B-7241 | 104.2 | 3.5 | 9.28 | 116.98 |
| | IMV Ac-5017 | 84.3 | – | 8.26 | 92.56 |
| Technical glycerol | IMV B-7241 | 122.0 | 363.9 | 7.36 | 493.26 |
| | IMV B-7405 | 139.9 | – | – | 139.9 |
| <i>n</i> -Hexadecane | IMV Ac-5017 | 44.8 | 21.4 | 5.73 | 71.93 |
| Refined oil | IMV B-7241 | 39.6 | 75.1 | 8.0 | 122.7 |
| | IMV B-7405 | 770.4 | 348.0 | 5.96 | 1124.36 |
| | IMV Ac-5017 | 19.4 | 17.1 | 7.8 | 44.3 |
| Waste oil after frying meat | IMV B-7241 | 83.2 | 43.6 | 9.49 | 136.29 |
| | IMV B-7405 | 23.3 | 53.9 | 46.8 | 124.0 |
| | IMV Ac-5017 | 91.3 | 37.8 | 5.83 | 134.93 |
| Waste oil after frying potatoes | IMV B-7405 | 84.7 | 15.9 | 18.16 | 118.76 |

Despite the fact that under the cultivation on such substrates as refined oil and glycerol, the total concentration of the phytohormones (500-1100 µg·L⁻¹) is the highest, their use for obtaining microbial preparations is limited due to the high cost. That's why the synthesis of phytohormones on the waste oil deserves attention: for all strains, the total concentration was in the range of 120-140 µg·L⁻¹. This level of synthesis was lower compared with the microorganisms, which are traditionally used as the growth

stimulators of plant (they synthesize from 600 to 40000 $\mu\text{g}\cdot\text{L}^{-1}$ phytohormones) [18, 19, 21]. But, given that the phytohormones show their stimulating effect in the extremely low concentrations ($10^{-5} - 10^{-12}$ $\text{mol}\cdot\text{L}^{-1}$), the rates of their synthesis by the producers of surfactants is acceptable for practical use in the plant growing.

Our previous research has shown that the surfactants of *N. vaccinii* IMV B-7405, *A. calcoaceticus* IMV B-7241 and *R. erythropolis* IMV Ac-5017 are the preparations of multifunctional purpose. So, they intensify the processes of decomposition of oil in water and soil, they have the antimicrobial activity, including against phytopathogenic bacteria; and also the surfactants of studying strains are characterized with the anti-adhesive properties and the ability to destroy biofilms [36 – 40].

Compared with the other microorganisms that are able to synthesize a complex of biologically active substances, in particular, phytohormones and metabolites with antimicrobial, nematocidal etc. activity [9 – 15], the complex preparations of strains *N. vaccinii* IMV B-7405, *A. calcoaceticus* IMV B-7241 and *R. erythropolis* IMV Ac-5017 have the following advantages.

First, these preparations have much wider range of properties (besides the antimicrobial and growth stimulating activities, also the destruction of oil, the anti-adhesive properties and the ability to destroy biofilms). Due to that they can be used not only in the plant growing for growth stimulation and control of pests number, but also in the environmental technologies for the environmental purification, in food, in pharmaceutical industry and medicine as the antimicrobial and antiadhesive agents.

Second, according to the literature data the antimicrobial metabolites synthesized simultaneously with the phytohormones are mostly characterized by an antifungal (rarely – nematocidal) activity. If it is antibacterial activity, then such products of microbial metabolism are antibiotics. Thus, streptomycetes synthesize geldanamycin [11], avermectins [13], blasticidin S, kasugamycin, oligomycin A, paramycin and pyrroles [14], and therefore, their use may be accompanied by the emergence of resistant forms of microorganisms. The mechanism of antimicrobial activity of surfactants is to ruin the integrity of the cytoplasmic membrane of the test cultures, which leads to the loss of cell viability [41]. In this case, the appearance of the microorganisms resistant to the microbial surfactants is unlikely.

Third, the microorganisms synthesize phytohormones on the rich nutrient mediums, which contain glucose, sucrose, dextrose, glucuronic acid, peptone, tryptone, mannitol as the source of carbon, and also tryptophan is being added exogenously to the medium for auxin formation, as it is the precursor of biosynthesis [42, 43]. Our research have shown the possibility of formation of the phytohormones of auxin, cytokinin [7] and gibberellin nature on the cheap mediums using toxic industrial waste (in particular, the waste oil and the technical glycerol as the waste of biodiesel production) as the substrates, as well as the possibility of the auxin synthesis without adding tryptophan to the medium [7].

Thus, the implementation of technologies based on *N. vaccinii* IMV B-7405, *A. calcoaceticus* IMV B-7241 and *R. erythropolis* IMV Ac-5017 will allow us not only to dispose the toxic industrial waste, but also to obtain the complex microbial preparations of wide range of application in the one process.

CONCLUSIONS

The results obtained earlier and presented in this article are the basis for the development of the waste-free technology using *N. vaccinii* IMV B-7405, *A. calcoaceticus* IMV B-7241 and *R. erythropolis* IMV Ac-5017 that will allow obtaining in a single process the microbial preparations with the various biological properties. Thus, when receiving surfactants, the precipitated cells can be used to purify water from oil; the obtained supernatant of the culture broth - for further separation of the surfactants with anti-adhesive and antimicrobial properties (including against the phytopathogenic bacteria). Since the aqueous phase, which remains after extraction of the surfactants, contains the phytohormones of auxin, cytokinin and gibberellic nature, it can be used to stimulate the growth of the microorganisms and plants.

REFERENCES

- Colla, L.M., Rizzardì, J., Pinto, M. H., Reinehr, C.O., Bertolin, T.E., Costa, J.A. V.: Simultaneous production of lipases and biosurfactants by submerged and solid-state bioprocesses, *Bioresource technology*, **2010**, **101** (21), 8308-8314;
- Hori, K., Ichinohe, R., Unno, H., Marsudi, S.: Simultaneous syntheses of polyhydroxyalkanoates and rhamnolipids by *Pseudomonas aeruginosa* IFO3924 at various temperatures and from various fatty acids, *Biochemical Engineering Journal*, **2011**, **53** (2), 196-202;
- Liang, T.W., Wu, C.C., Cheng, W.T., Chen, Y.C., Wang, C.L., Wang, I.L., Wang, S.L.: Exopolysaccharides and antimicrobial biosurfactants produced by *Paenibacillus macerans* TKU029, *Applied biochemistry and biotechnology*, **2014**, **172** (2), 933-950;
- Sharma, D., Singh Saharan, B.: Simultaneous production of biosurfactants and bacteriocins by probiotic *Lactobacillus casei* MRTL3, *International journal of microbiology*, **2014**, **2014**, 1-7;
- Bhange, K., Chaturvedi, V., Bhatt, R.: Simultaneous production of detergent stable keratinolytic protease, amylase and biosurfactant by *Bacillus subtilis* PF1 using agro industrial waste, *Biotechnology reports*, **2016**, **10**, 94-104;
- Zarinviarsagh, M., Ebrahimipour, G., Sadeghi, H.: Lipase and biosurfactant from *Ochrobactrum* intermedium strain MZV101 isolated by washing powder for detergent application, *Lipids in health and disease*, **2017**, **16** (1), 177;
- Pirog, T., Leonova, N., Shevchuk, T., Savenko, I., Iutinska, H.: [Synthesis of phytohormones bacteria of *Acinetobacter calcoaceticus* IMV B-7241, *Rhodococcus erythropolis* IMV Ac-5017 and *Nocardia vaccinii* IMV B-7405 – producers of surface-active substances]: *Proceedings of the National Academy of Sciences of Belarus. Biological series* (Editors: Nikiforov M.Y., Volotovskiy I.D., Evtushenkov A.N., **2016**, 90–95 (In Russian);
- Pacwa-Płociniczak, M., Płociniczak, T., Iwan, J., Żarska, M., Chorążewski, M., Dzida, M., Piotrowska-Seget, Z.: Isolation of hydrocarbon-degrading and biosurfactant-producing bacteria and assessment their plant growth-promoting trait, *Journal of environmental management*, **2016**, **168**, 175-184;
- Grady, E.N., MacDonald, J., Liu, L., Richman, A., Yuan, Z.: Current knowledge and perspectives of *Paenibacillus*: a review, *Microbial cell factories*, **2016**, **15** (1), 203;
- Kaur, T., Manhas, R.: Antifungal, insecticidal, and plant growth promoting potential of *Streptomyces hydrogenans* DH16, *Journal of basic microbiology*, **2014**, **54** (11), 1175-1185;
- Boudjeko, T., Tchinda, R.A.M., Zitouni, M., Nana, J.A., Lerat, S., Beaulieu, C.: *Streptomyces cameroonensis* sp. nov., a geldanamycin producer that promotes *Theobroma cacao* growth, *Microbes and environments*, **2017**, ME16095;
- Matsumoto, A., Takahashi, Y.: Endophytic actinomycetes: promising source of novel bioactive compounds, *The Journal of Antibiotics*, **2017**, **70** (5), 514;
- Iutynska, G.O., Biliavska, L.O., Kozyriska, V.Y.: Development strategy for the new environmentally friendly multifunctional bioformulations based on soil streptomycetes, *Mikrobiologichnyi zhurnal*, **2017**, **79** (1), 22-33;

14. Law, J.W.F., Ser, H.L., Khan, T.M., Chuah, L.H., Pusparajah, P., Chan, K.G., Lee, L.H.: The potential of *Streptomyces* as biocontrol agents against the rice blast fungus, *Magnaporthe oryzae* (*Pyricularia oryzae*), *Frontiers in microbiology*, **2017**, 8, 3;
15. Ali, S., Hameed, S., Imran, A., Iqbal, M., Lazarovits, G.: Genetic, physiological and biochemical characterization of *Bacillus* sp. strain RMB7 exhibiting plant growth promoting and broad spectrum antifungal activities, *Microbial cell factories*, **2014**, 13 (1), 144;
16. Tsavkelova, E.A., Klimova, S.Y., Cherdynitseva, T.A., Netrusov, A.I.: Microbial producers of plant growth stimulators and their practical use: a review, *Applied biochemistry and microbiology*, **2016**, 42 (2), 117-126;
17. Singh, S.: A review on possible elicitor molecules of cyanobacteria: their role in improving plant growth and providing tolerance against biotic or abiotic stress, *Journal of applied microbiology*, **2014**, 117 (5), 1221-1244;
18. Vejan, P., Abdullah, R., Khadiran, T., Ismail, S., Nasrulhaq Boyce, A.: Role of plant growth promoting rhizobacteria in agricultural sustainability - a review, *Molecules*, **2016**, 21 (5), 573;
19. Radhakrishnan, R., Hashem, A., Abd Allah, E.F.: *Bacillus*: a biological tool for crop improvement through bio-molecular changes in adverse environments, *Frontiers in physiology*, **2017**, 8, 667;
20. Pirog, T.P., Shevchuk, T.A., Voloshina, I.N., Gregirchak, N.N.: Use of claydite-immobilized oil-oxidizing microbial cells for purification of water from oil, *Applied Biochemistry and Microbiology*, **2005**, 41 (1), 51-55;
21. Muromtsev, G.S., Agnistikova, V.N.: *Gibberellins plant hormones*, Nauka, Moscow, **1973** (in Russian);
22. Lakin, G.F.: *Biometriya (Biometry)*, Vysshaya shkola, Moscow, **1990** (in Russian);
23. Hedden, P.: *Annual plant reviews, the gibberellins*, Vol. 49, Thomas GS, eds., Wiley Blackwell, NJ, **2016**;
24. Muddapur, U.M., Gadkari, M.V., Kulkarni, S.M., Sabannavar, P.G., Niyonzima, F.N., More, S.S.: Isolation and characterization of gibberellic acid 3 producing *Fusarium* sp. from Belgaum agriculture land Andits impact on green pea and rice growth promotion, *Aperito journal of advanced plant biology*, **2015**, 1 (2), 106;
25. Khan, A.L., Hussain, J., Al-Harrasi, A., Al-Rawahi, A., Lee, I.J.: Endophytic fungi: resource for gibberellins and crop abiotic stress resistance, *Critical reviews in biotechnology*, **2015**, 35 (1), 62-74;
26. Leitão, A.L., Enguita, F.J.: Gibberellins in *Penicillium* strains: challenges for endophyte-plant host interactions under salinity stress, *Microbiological research*, **2016**, 183, 8-18;
27. Jaroszek-Ścisiel, J., Kurek, E., Trytek, M.: Efficiency of indoleacetic acid, gibberellic acid and ethylene synthesized in vitro by *Fusarium culmorum* strains with different effects on cereal growth, *Biologia*, **2014**, 69 (3), 281-292;
28. Shi, T.Q., Peng, H., Zeng, S.Y., Ji, R.Y., Shi, K., Huang, H., Ji, X.J.: Microbial production of plant hormones: opportunities and challenges, *Bioengineered*, **2017**, 8 (2), 124-128;
29. Ma, K.W., Ma, W.: Phytohormone pathways as targets of pathogens to facilitate infection, *Plant molecular biology*, **2016**, 91 (6), 713-725;
30. Rashad, F.M., Fathy, H.M., El-Zayat, A.S., Elghonaimy, A.M.: Isolation and characterization of multifunctional *Streptomyces* species with antimicrobial, nematocidal and phytohormone activities from marine environments in Egypt, *Microbiological research*, **2015**, 175, 34-47;
31. Atzorn, R., Crozier, A., Wheeler, C.T., Sandberg, G.: Production of gibberellins and indole-3-acetic acid by *Rhizobium phaseoli* in relation to nodulation of *Phaseolus vulgaris* roots, *Planta*, **1988**, 175 (4), 532-538;
32. Makeeva, A.: Research of microbial synthesis individual gibberellins and abscisic acid, Dissertation, All-Russia Research Institute of Agricultural Biotechnology, **1996** (In Russian);
33. Park, Y.G., Mun, B.G., Kang, S.M., Hussain, A., Shahzad, R., Seo, C.W., Lee, I.J.: *Bacillus aryabhattai* SRB02 tolerates oxidative and nitrosative stress and promotes the growth of soybean by modulating the production of phytohormones, *PLoS One*, **2017**, 12 (3), e0173203;
34. Khan, A.L., Waqas, M., Kang, S.M., Al-Harrasi, A., Hussain, J., Al-Rawahi, A., Lee, I.J.: Bacterial endophyte *Sphingomonas* sp. LK11 produces gibberellins and IAA and promotes tomato plant growth, *Journal of Microbiology*, **2014**, 52 (8), 689-695;
35. Kang, S.M., Khan, A.L., You, Y.H., Kim, J.G., Kamran, M., Lee, I.J.: Gibberellin production by newly isolated strain *Leifsonia soli* SE134 and its potential to promote plant growth, *Journal of microbiology and biotechnology*, **2014**, 24 (1), 106-112;

36. Pirog, T., Sofilkanych, A., Shevchuk, T., Shulyakova, M.: Biosurfactants of *Rhodococcus erythropolis* IMV Ac-5017: synthesis intensification and practical application, *Applied biochemistry and biotechnology*, **2013**, 170 (4), 880-894;
37. Pirog, T., Sofilkanych, A., Konon, A., Shevchuk, T., Ivanov, S.: Intensification of surfactants' synthesis by *Rhodococcus erythropolis* IMV Ac-5017, *Acinetobacter calcoaceticus* IMV B-7241 and *Nocardia vaccinii* K-8 on fried oil and glycerol containing medium, *Food and bioproducts processing*, **2013**, 91 (2), 149-157;
38. Pirog, T.P., Konon, A.D., Sofilkanich, A.P., Iutinskaya, G.A.: Effect of surface-active substances of *Acinetobacter calcoaceticus* IMV B-7241, *Rhodococcus erythropolis* IMV Ac-5017, and *Nocardia vaccinii* K-8 on phytopathogenic bacteria, *Applied biochemistry and microbiology*, **2013**, 49 (4), 360-367;
39. Pirog, T.P., Konon, A.D., Beregovaya, K.A., Shulyakova, M.A.: Antiadhesive properties of the surfactants of *Acinetobacter calcoaceticus* IMB B-7241, *Rhodococcus erythropolis* IMB Ac-5017, and *Nocardia vaccinii* IMB B-7405, *Microbiology*, **2014**, 83 (6), 732-739;
40. Pirog, T.P., Panasyuk, E.V., Antonyuk, N.A. Impact of microbial *Nocardia vaccinii* IMB B-7405 surfactants on oil destruction in water, *Journal of water chemistry and technology*, **2016**, 38 (5), 301-306;
41. Rodrigues, L., Banat, I.M., Teixeira, J., Oliveira, R.: Biosurfactants: potential applications in medicine. *Journal of antimicrobial chemotherapy*, **2006**, 57 (4), 609-618;
42. Majeed, A., Abbasi, M.K., Hameed, S., Imran, A., Rahim, N.: Isolation and characterization of plant growth-promoting rhizobacteria from wheat rhizosphere and their effect on plant growth promotion, *Frontiers in microbiology*, **2015**, 6, 198;
43. Giassi, V., Kiritani, C., Kupper, K.C.: Bacteria as growth-promoting agents for citrus rootstocks, *Microbiological research*, **2016**, 190, 46-54.